arXiv:1205.3195v1 [quant-ph] 14 May 2012

Decoherence and the Quantum Detection of Classically Undetectable Phenomena

C. Jess Riedel
Theoretical Division / CNLS, LANL, Los Alamos, New Mexico 87545, USA and
Department of Physics, University of California, Santa Barbara, California 93106, USA

(Dated: March 4, 2013)

I consider the general problem of exploiting quantum mechanics to detect weak particles and forces. A phenomenon is detectable using classical methods if and only if the effective Hamiltonian coupling it to normal matter does not commute with the classically preferred basis. Classically undetectable phenomena can still be observed, but they require coherent superpositions and measurements which do not commute with the preferred basis. The Aharonov-Bohm effect, the shot noise limit, and the standard quantum limit are special cases of this general idea. I propose using matter-wave interferometry to directly detect hypothetical new particles, such as sub-MeV dark matter, that transfer negligible momentum when colliding with normal matter. The coherent control of planck-mass superpositions could also enable the detection of relic neutrinos, fifth forces, and even gravitons.

One limit of experimental physics is that it does not allow us to rule out the existence of new particles, forces, dimensions, or universes from which we are causally disconnected. Even if some hypothetical new physics has non-zero interactions with experimental apparatuses, the coupling may always be so small that the influence on the apparatuses is negligible. Experimenters are only able to rule out coupling strengths above some cutoff, and at some level of statistical significance. There are many proposed and underway experimental searches for hypothetical new particles and forces (with varying degrees of theoretical and observational motivation) which, if they exist, interact only weakly with regular matter. Most notable is probably dark matter [1–5], but others include supersymmetric particles [6], new neutrino flavors [7], mirror matter [8], and fifth forces [9].

Methods for detecting weak phenomena can often be described by the following classical detection scheme. There is a normal system $\mathcal{A}$, which is relatively easy to handle and measure, and there is an environment $\mathcal{E}$, which continuously interacts with $\mathcal{A}$. There is also a hypothetical system $\mathcal{D}$ which is difficult to measure but interacts at least weakly with $\mathcal{A}$. The coupling between $\mathcal{A}$ and $\mathcal{E}$ determines a (possibly overcomplete) preferred basis $\{|A_\alpha\rangle\}$ in $\mathcal{A}$ of *pointer* states [10, 11]. (For example, $\mathcal{A}$ is a Xenon nucleus, $\mathcal{E}$ is the rest of a bath of liquid Xenon, $\mathcal{D}$ is a dark matter candidate, and the $\{|A_\alpha\rangle\}$ are wavepackets [12] localized in phase space.) Decoherence ensures that the state of $\mathcal{A}$ is approximately diagonal in the preferred basis, both before and after interacting with $\mathcal{D}$. Schematically: the initial density matrix

$$\rho^0_{\mathcal{A}} = \sum_i p_i |A_\alpha\rangle\langle A_\alpha| \tag{1}$$

of $\mathcal{A}$ undergoes some unitary evolution,

$$\rho^0_{\mathcal{A}} \otimes \rho^0_{\mathcal{D}} \to \rho_{\mathcal{A}\mathcal{D}}, \tag{2}$$

and is then measured in the preferred basis $\{|A_\alpha\rangle\}$. Even when there is not a natural environment enforcing a pointer basis, there is often still a single preferred basis in which the system is prepared and in which it can be measured, due to experimental limitations.

Besides the advantages of conceptual simplicity, searches for weak phenomena commonly employ a classical detection scheme because weak phenomena are often *rare* phenomena, so a reasonable chance of observations necessitates large experimental volumes. It's usually not possible to prepare large volumes in arbitrary quantum states, nor make arbitrary quantum measurements. Indeed, the initial state $\rho^0_{\mathcal{A}}$ of $\mathcal{A}$ will be approximately diagonal in the basis $\{|A_\alpha\rangle\}$ because of decoherence [13–15] from the environment $\mathcal{E}$. Furthermore, an experimenter's instruments do not interact with $\mathcal{A}$ directly. Instead, he uses the environment $\mathcal{E}$ as an information channel [16, 17], and this turns out to prevent him from measuring $\mathcal{A}$ in anything besides the pointer basis [18].

Crucially, such detection schemes rely on the hypothetical system changing the state of the regular matter *in the preferred basis*. Let $\rho_{\mathcal{A}}$ and $\tilde{\rho}_{\mathcal{A}}$ be the state of $\mathcal{A}$ before measurement if $\mathcal{D}$ does or does not exist, respectively. A necessary condition for the detection of $\mathcal{D}$ is for

$$(\rho_{\mathcal{A}})_{\alpha\alpha} = \langle A_\alpha|\rho_{\mathcal{A}}|A_\alpha\rangle \tag{3}$$

to be substantially different than $(\tilde{\rho}_{\mathcal{A}})_{\alpha\alpha}$ for at least one value of α.

For example, many experiments consist essentially of a container of regular matter (e.g. liquid hydrocarbons [7, 19], heavy water [20], nobel liquids [21], or crystaline semiconductors [4]) which is carefully watched for the tiny effects of a collision with the hypothetical particle such as recoil, vibration, heating, or ionization. These techniques rely on there being sufficient momentum transfer from the hypothetical particle to the regular matter that the momentum of the regular matter is substantially changed.

But it's easy to imagine hypothetical new particles which transfer negligible momentum. Classically, such particles are *undetectable*. In a classical universe, or a quantum universe where experimenters only employ classical detection schemes, the new particles are ghosts. They evolve along side normal matter but cannot influence the normal matter even though the normal matter can (strongly) influence them. Their existence may be motivated only by elegance, or perhaps (like dark matter) by their astrophysical and cosmological influence through gravity.

More generally, a classical detection scheme will be blind to new phenomena (for all possible initial states) if and only if the unitary governing the joint evolution is diagonal in the preferred basis. To see this, first suppose that the evolution is of the form

$$U|A_i\rangle|D_\alpha\rangle = |A_i\rangle U^{(i)}|D_\alpha\rangle, \tag{4}$$

where U acts on $\mathcal{A}\otimes\mathcal{D}$ and the $U^{(i)}$ act on $\mathcal{D}$. (This is equivalent to U commuting with $\{|A_i\rangle\}$, i.e. all eigenvalues of U are of the form $|A_i\rangle|D^{(i)}_\beta\rangle$.) Then the matrix elements of the state of $\mathcal{A}$ in the preferred basis after the interaction are

$$\langle A_i|\rho_\mathcal{A}|A_j\rangle = \langle A_i|\,\mathrm{Tr}_\mathcal{D}\left[U\left(\rho^0_\mathcal{A}\otimes\rho^0_\mathcal{D}\right)U^\dagger\right]|A_j\rangle \tag{5}$$

$$= \langle A_i|\rho^0_\mathcal{A}|A_j\rangle\,\mathrm{Tr}_\mathcal{D}\left[U^{(i)}\rho^0_\mathcal{D}U^{(j)\dagger}\right] \tag{6}$$

Therefore, $\rho_\mathcal{A}$ can differ from the initial state $\rho^0_\mathcal{A}$ *only* on the off-diagonal elements (necessitating a measurement outside the preferred basis) and *only* if $\rho^0_\mathcal{A}$ itself has non-zero off-diagonal elements (necessitating an initial coherent superposition of preferred states).

Conversely, suppose that $\rho^0_\mathcal{A}$ is diagonal in the preferred basis and $\mathcal{D}$ is in arbitrary pure initial state $\rho^0_\mathcal{D} = |D_0\rangle\langle D_0|$. Then the diagonal elements of the state of $\mathcal{A}$ are

$$\langle A_k|\rho_\mathcal{A}|A_k\rangle = \sum_i\langle A_i|\rho^0_\mathcal{A}|A_i\rangle\langle D_0^{(k,i)}|D_0^{(k,i)}\rangle, \tag{7}$$

where $|D_0^{(k,i)}\rangle = \langle A_k|U|A_i\rangle|D_0\rangle$. Blindness of the classical detection scheme implies that $\langle D_0^{(k,i)}|D_0^{(k,i)}\rangle = \delta_{k,i}$ for all $|D_0\rangle$, which can only be true if Eq. (4) holds. Furthermore, the previous paragraphs show that relaxing either restriction of the classical detection scheme (that is, allowing either initial states or measurements outside the preferred basis) will not lead to a visibility of the phenomena; if it can't be seen with a classical detection scheme, it can only be seen if both the initial state and the measurement fail to commute with the preferred basis.

In the case of hypothetical particles with negligible momentum, scattering events (which we assume happen on much shorter timescale than the free propagation of the regular matter) do not substantially influence a localized wavepacket of the regular matter:

$$U_\mathrm{scatt}|A_{\vec{x},\vec{p}}\rangle|D_\mathrm{in}\rangle \approx |A_{\vec{x},\vec{p}}\rangle|D_\mathrm{out}\rangle, \tag{8}$$

where $|A_{\vec{x},\vec{p}}\rangle$ is a wavepacket of $\mathcal{A}$ at position $\vec{x}$ with momentum $\vec{p}$.

Nevertheless, such particles are detectable, maybe even easily detectable, with specifically quantum experiments. Before I discuss this, I turn to the Aharonov-Bohm effect as a prototypical example of the power of non-classical detection schemes.

The Aharonov-Bohm effect

The startling Aharonov-Bohm (AB) effect [22, 23] has been experimentally demonstrated many times, in both magnetic [24–26] and electric [27] versions. Charged particles are made to travel coherently along two alternate paths, such as in a Mach-Zehnder interferometer. At the experimenter's discretion, an external electromagnetic potential can be applied so that the two paths are at different potentials yet still experience zero magnetic and electric field. The paths are recombined, and the size of the potential difference determines the phase of the interference pattern. The effect is often interpreted as a demonstration that the electromagnetic potential is physically 'real', rather than just a useful mathematical concept.

However, Vaidman recently pointed out that this is a mistaken interpretation which is an artifact of the semi-classical approximation used to describe the AB effect [28]. Although it is true that the superposed test charges experience zero field, it turns out that the *source* charges creating that macroscopic potential *do* experience a non-zero field, and that the strength of this field is dependent on which path is taken by the test charges. In fact, if one modifies the experiment so that the source charges experience zero field, while retaining a potential difference between the two paths, the AB effect disappears. (In this modification, the semi-classical approximation breaks down.) It is the non-zero field at the source charges—*not* the potential at the test charges—which is responsible for the observed phase.

Even though the AB effect doesn't tell us anything new about the electromagnetic potential, it turns out to be a prototypical example of using a non-classical detection scheme to detect a classically undetectable phenomenon. Classically, there is no way to distinguish between two electromagnetic field configurations by performing experiments in a region which always has zero field. This is equivalent to saying that the unitary used to evolve the state of the test charges commutes with the pointer basis of test charge wavepackets localized in phase space, $\{|A_{\vec{x},\vec{p}}\rangle\}$, if the field is zero for all relevant values of $\vec{x}$. Indeed, suppose $|D_\mathrm{off}\rangle$ features zero electromagnetic potential everywhere and $|D_\mathrm{on}\rangle$ features a spatial region R with uniform electric potential V_0. Consider two pointer states $|A_{\vec{x},0}\rangle$ and $|A_{\vec{y},0}\rangle$, with $\vec{y}\in R$ but $\vec{x}\notin R$, for a test


$|A_{\vec{x},\vec{p}}\rangle$
$|D^{(\vec{x})}_{\text{out}}\rangle$
$\mathcal{A}$
$|A_{\vec{y},\vec{p}}\rangle$
$|D_{\text{in}}\rangle$
$\mathcal{D}$
$|D^{(\vec{y})}_{\text{out}}\rangle$

FIG. 1. Decoherence detection with a Mach-Zehnder Interferometer. System $\mathcal{A}$ is places in a coherent superposition of spatially displaced wavepackets by a beam splitter which each travel a separate path and then are recombined. In the absence of system $\mathcal{D}$, the interferometer is tuned so that $\mathcal{A}$ will be detected at the bright port with near unit probability, and at the dark port with near vanishing probability. However, if system $\mathcal{D}$ scatters off $\mathcal{A}$, these two paths will decohere and $\mathcal{A}$ will be detected at either port with equal probability.

charge $-e$. The effective Hamiltonian is

$$H = -eV_0 |A_{\vec{y},0}\rangle\langle A_{\vec{y},0}| \otimes |D_{\text{on}}\rangle\langle D_{\text{on}}|, \tag{9}$$

which clearly commutes with the pointer basis $\{|A_{\vec{x},0}\rangle, |A_{\vec{y},0}\rangle\}$. That is, after a time T the joint state picks up a phase $e^{i\phi}$, where $\phi = -eV_0T$ if the test charge is in region R and the macroscopic electromagnetic potent is switched on, and $\phi = 0$ otherwise.

A classical detection scheme in the pointer basis cannot distinguish whether the macroscopic field corresponds to $|D_{\text{on}}\rangle$ or $|D_{\text{off}}\rangle$; the global phase is not observable. However, if the experimenter can place the test charge in a coherent superposition of $|A_{\vec{x},0}\rangle$ and $|A_{\vec{y},0}\rangle$ and later measure the test charge in a basis which does not commute with the pointer basis, he can determine the state of $\mathcal{D}$. Producing such a superposition and measurement is exactly the feature of AB-effect experiments that enables this classically impossible feat.

Detection of particles transferring negligible momentum

Returning to the problem of detecting hypothetical particles which transfer negligible momentum, it's now not hard to guess how to proceed: the experimenter must transcend the classical detection scheme by preparing $\mathcal{A}$ in a superposition of pointer states and measuring $\mathcal{A}$ in a basis which does not commute with the pointer basis. A natural choice is atom interferometry or, more generally, matter-wave interferometry [29, 30].

An atom interferometer takes advantage of the deBroglie wave nature of matter. For now, let us specialize to a Mach-Zehnder interferometer, Fig. 1, although the ideas will apply generally. An atom is prepared in a coherent superposition $|A_{\vec{x},\vec{p}}\rangle + |A_{\vec{y},\vec{p}}\rangle$ of two wavepackets with something analogous to a beam splitter. These wavepackets are allowed to propagate over some length, and then the wavepackets are recombined with a second splitter. Assuming the spread of the wavepackets is negligible, we can align the second splitter so that the atom ends up at one "bright" detector with near unit probability, and at the other "dark" detector with near vanishing probability.

The detectors effectively measure $\mathcal{A}$ in the basis $\{|A_\pm\rangle = |A_{\vec{x},\vec{p}}\rangle \pm |A_{\vec{y},\vec{p}}\rangle\}$. If we let the state $|D_\varnothing\rangle$ represent the absence of the hypothetical particles, we can take the evolution to be trivial:

$$\Big[|A_{\vec{x},\vec{p}}\rangle + |A_{\vec{y},\vec{p}}\rangle\Big]|D_\varnothing\rangle \to \Big[|A_{\vec{x},\vec{p}}\rangle + |A_{\vec{y},\vec{p}}\rangle\Big]|D_\varnothing\rangle, \tag{10}$$

Measuring in the basis $\{|A_\pm\rangle\}$ gives outcome $|A_+\rangle$ with certainty. But suppose a hypothetical particle decoheres the superposition by scattering off the atom, thereby recording which-path information,

$$\Big[|A_{\vec{x},\vec{p}}\rangle + |A_{\vec{y},\vec{p}}\rangle\Big]|D_{\text{in}}\rangle \to |A_{\vec{x},\vec{p}}\rangle|D^{(\vec{x})}_{\text{out}}\rangle + |A_{\vec{y},\vec{p}}\rangle|D^{(\vec{y})}_{\text{out}}\rangle, \tag{11}$$

where $\langle D^{(\vec{x})}_{\text{out}}|D^{(\vec{y})}_{\text{out}}\rangle \approx 0$. In this case, a measurement in the basis $\{|A_\pm\rangle\}$ gives equal probability of either outcome. When the dark detector clicks (which it will do half the time), this gives direct evidence of the hypothetical particle $\mathcal{D}$ *even if it transfers negligible momentum to* $\mathcal{A}$. Importantly, the effective unitary governing the interaction,

$$U = I_{\mathcal{A}} \otimes |D_\varnothing\rangle\langle D_\varnothing| \tag{12}$$

$$+|A_{\vec{x},\vec{p}}\rangle\langle A_{\vec{x},\vec{p}}| \otimes \Big[|D^{(\vec{x})}_{\text{out}}\rangle\langle D_{\text{in}}| + \text{h.c.}\Big] \tag{13}$$

$$+|A_{\vec{y},\vec{p}}\rangle\langle A_{\vec{y},\vec{p}}| \otimes \Big[|D^{(\vec{y})}_{\text{out}}\rangle\langle D_{\text{in}}| + \text{h.c.}\Big], \tag{14}$$

is obviously diagonal in the basis $\{|A_{\vec{x},\vec{p}}\rangle, |A_{\vec{y},\vec{p}}\rangle\}$. There is no way to detect the presence of $\mathcal{D}$ using a classical detection scheme, but atom interferometry can expose it simply.

This basic idea is implied by the decoherence experiments of Hornberger et al. [31] and others [32]. In the Hornberger et al. experiments, coherent spatial superpositions of C_{70} fullerenes were demonstrated by passing them through several gratings and recording the interference pattern. (This is a generalized double-slit experiment rather than a Mach-Zehnder interferometer.) The interference region was filled with a gas of molecules much smaller than the fullerenes, such that collisions recorded which-way information in the gas but did not substantial deflect the fullerenes. The pressure of the gas was adjustable, so that (by observing the suppression of the interference fringes) the presence of the gas—and moreover its density—could be inferred even though the gas imparted negligible momentum to the measured system.

Of course, the imaginary atom-interferometry $\mathcal{D}$-detector only works if the flux of hypothetical $\mathcal{D}$ particles is high enough such that at least one will scatter off a $\mathcal{A}$ wavepacket with significant probability. More generally, the initial state of $\mathcal{D}$ will be described by some density matrix $\rho^0_{\mathcal{D}} = \sum_i d_i |D_i\rangle\langle D_i|$, and there will be conditional unitaries $U^{(\vec{x})}$ and $U^{(\vec{y})}$. Decoherence will only be effective if the average overlap

$$\sum_i d_i \langle D_i^{(\vec{x})} | D_i^{(\vec{y})} \rangle = \sum_i d_i \langle D_i | U^{(\vec{x})\dagger} U^{(\vec{y})} | D_i \rangle \tag{15}$$

is much less than unity.

There are at least two ways to increase the likelihood of a scattering event, and hence increase the sensitivity of the experiment to the hypothetical $\mathcal{D}$ particles. First, the time over which the superposition is maintained can be increased, possibly by increasing the length of the interferometer arms. For an uncorrelated $\mathcal{D}$ gas, the expected number of scattering events should be linear in time. Second, one can superpose ever larger clusters of atoms. That is, construct a matter-wave interferometer with as large of objects $\mathcal{A}$ as possible. As more atoms are added, the chance that at least one $\mathcal{D}$ particle scatters goes up, and it only takes a *single* scattering event to decohere the entire $\mathcal{A}$ system.

Of course, there are many possible sources of decoherence; anomalous decoherence hardly implies the existence of new particles. Still, note that the inverse statement *is* true: the observation of interference effects (which establishes the existence of the superposition, and hence implies that all relevant sources of decoherence have been eliminated) implies that there are *not* new particles (up to some level of sensitivity). Furthermore, if anomalous decoherence is observed, one can convincingly demonstrate that it is due to hypothetical particles by observing the functional dependence of the degree of decoherence on parameters which effect the scattering rate of the hypothetical particles. This is discussed further in the next section for the special case of dark matter searches.

Nimmrichter et al. have recently proposed [33] using matter-wave interferometers to test "objective collapse" theories, such as the theory of Ghirard, Rimini, and Weber (GRW) [34], which explicitly modify the Schrödinger equation to produce localized wavepackets stochastically. As Nimmrichter et al. point out, Vacchini has shown [35] that the master equation for the density matrix of a mass under GRW is equivalent to the master equation for that same mass when subjected to an appropriate type of collisional decoherence. This suggest caution in interpreting a potential observation of anomalous suppression of interference fringes as evidence of objective collapse; the suppression might be explained by uncontrolled collisional decoherence described by normal quantum mechanics.

If experimenters only had to worry about known known interactions, this would not be too worrying because gas pressure, ambient light, and blackbody radiation can be well understood. However, heretofore unknown particles and forces have the potential to mimic objective collapse in state-of-the-art experiments.

Dark matter direct detection through matter-wave interferometry

Astrophysical and cosmological evidence suggest that the majority of mass in the universe is made up of invisible dark matter. Existing experiments, observations, and theoretical preferences provide a complicated thicket of conditional restrictions on any potential theory of dark matter, but model-independent constraints are still rare and very valuable. Here I show that the methods described in the last section are sensitive to types of dark matter which have not been accessed by conventional direct-detection techniques. I will concentrate on the popular cold dark matter scenario which is supported by a bare minimum of assumptions.

Based just on the movement of luminous matter in the galaxy and the virial theorem, we infer that cold dark matter in our region of the Milky Way has a density of roughly $\rho \sim 0.4$ GeV/cm^3 [36] and a (non-relativistic) Maxwellian velocity distribution centered around $\bar{v} \sim 230$ km/s with a cutoff at the galactic escape velocity $v_{\mathrm{e}} \sim 525$ km/s [37]. For now, I assume that there is no mechanism that would substantially change ρ in the immediate vicinity of the Earth (though see, for example, [38]). Given this, any hypothetical mass m_{DM} for the dark matter particle then fixes the rough interaction rate of a particle of normal matter (either an electron or a nucleon) as a function of scattering cross section: $\Gamma_0 \approx v\sigma\rho/m_{\mathrm{DM}}$. This is why the most popular parameterization of the cold dark matter theory space is in terms of the mass m_{DM} and a cross-section σ. Typically, this is the total cross section for spin-independent scattering off nucleons.

Direct detection experiments look for signs of recoil when a dark matter particle collides with normal matter of mass M, usually elastically. The energy deposited is $\mathrm{O}(m_{\mathrm{DM}}^2 v^2/M)$. So long as this recoil is measurable, it is doubtlessly easiest to use a classical detection scheme for the reason stated in the introductory section: nonclassical states of matter are difficult to prepare and measure, and this problem is exacerbated by the large volumes necessary to observe rare events.

For nuclear recoil, direct detection experiments are sensitive down to a few keV being deposited in the detector, which corresponds to a sensitivity to dark matter masses greater than a few GeV. This lines up well with the Lee-Weinberg bound [39], which constrains the most popular and natural form of WIMP dark matter to a mass of at least 2 GeV.

However, the most natural WIMP models have seen their parameter space severely constricted by experimental searches [] and astrophysical observations []. It is prudent to allow for more general possibilities, and there are several models of cold dark matter with sub-GeV mass [40–45]. The mass range between an MeV and a GeV is at least testable by looking at electron, rather than nuclear, recoil [46] because the lighter electrons absorb more energy and also have a lower detectable energy threshold than nuclei. But there is no a priori relation between nucleon and electron scattering, and direct detection techniques which identify individual collisions probably cannot do better than this. A 1 MeV dark matter candidate will typically deposit less than an eV when colliding with an electron and less than 10^{-3} eV when colliding with a nucleus. (For a keV candidate, the values are 10^{-6} and 10^{-9} eV, respectively.) Therefore, sub-MeV dark matter is a promising scenario for decoherence-sensing interferometers to have a comparative advantage.

Consider a coherent spatial superposition of normal matter $|A_{\vec{x},\vec{p}}\rangle + |A_{\vec{y},\vec{p}}\rangle$, where the separation $\vec{\Delta x} = \vec{x} - \vec{y}$ is at least as large as the width of the wavepackets. The ability of dark matter to decohere this state is determined by the overlap $\langle D^{(\vec{x})}_{\text{out}} | D^{(\vec{y})}_{\text{out}} \rangle$ of the conditional dark-matter post-scattering states. This overlap, in turn, is strongly effected by size of the de Broglie wavelength $\lambda = 2\pi\hbar / v m_{\text{DM}}$ of the scattered dark matter particle compared to the separation Δx. If $\Delta x \ll \lambda$, then $|\langle D^{(\vec{x})}_{\text{out}} | D^{(\vec{y})}_{\text{out}} \rangle|$ is only slightly away from unity. The dark matter cannot easily "see" the superposition, and it takes many scattering events to decohere the normal matter [51]. On the other hand, if $\Delta x \gg \lambda$, then $\langle D^{(\vec{x})}_{\text{out}} | D^{(\vec{y})}_{\text{out}} \rangle$ nearly vanishes. The out states are distinct and a single dark matter collision suffices to decohere.

The general case was worked out by Gallis and Fleming [52]. (See also [53–55].) The state of the normal matter in the $\{|A_{\vec{x},\vec{p}}\rangle, |A_{\vec{y},\vec{p}}\rangle\}$ basis after a time T is

$$\rho_{\mathcal{A}} = \frac{1}{2} \begin{pmatrix} 1 & \gamma \\ \gamma^* & 1 \end{pmatrix} \tag{16}$$

where $\gamma = \exp[-T F(\vec{\Delta x})]$ is the decoherence factor. The decoherence rate $F(\vec{\Delta x})$ is given by

$$\begin{aligned} F(\vec{\Delta x}) = \int \mathrm{d}q\, n(q) v(q) \int \mathrm{d}\hat{q}\, \mathrm{d}\hat{q}' \frac{1}{2} \\ \times \left\{1 - \cos[q(\hat{q} - \hat{q}') \cdot \vec{\Delta x}]\right\} |f(q\hat{q}, q\hat{q}')|^2 \end{aligned} \tag{17}$$

where $n(q)$ is the spectral number density (number per unit volume, per unit momentum) of dark matter, $|f(q\hat{q}, q\hat{q}')|^2$ is the differential cross section, and $v(q)$ is the velocity of dark matter with momentum q. For cold dark matter, this is simply $v(q) = q/m_{\text{DM}}$. The angular integrals can only be done by assuming a form for the differential cross section, but the excluded region will not be very sensitive to it so long as it does not vary to fast with momentum q. In the small wavelength limit, F will be a function of the total cross section $\sigma^{\text{tot}} = \int\int \mathrm{d}\hat{q}\, \mathrm{d}\hat{q}' |f(q\hat{q}, q\hat{q}')|^2 / 4\pi$. In the large wavelength limit, F will be a function of the momentum-transfer cross section $\sigma^{\text{tr}} = \int\int \mathrm{d}\hat{q}\, \mathrm{d}\hat{q}' (1 - \cos\theta) |f(q\hat{q}, q\hat{q}')|^2 / 4\pi$, where θ is the scattering angle. For the purposes of plotting exclusion regions, I will take $|f(q\hat{q}, q\hat{q}')|^2 \propto (1 + \cos^2\theta)$, which is the form of Thomson and Rayleigh scattering. For this choice, and also for the choice of hard-sphere



FIG. 2. The dark matter detection reach of several proposed experiments featuring macroscopic spatial superpositions, compared with existing constraints, for the spin-independent nucleon scattering cross section. The dotted lines labeled 'Rocket', 'Balloon', and 'Sea level' denote maximum cross-sections accessible by sounding rockets, sounding balloons, and terrestrial experiments at their respective altitudes due to atmospheric shielding. The X-ray Quantum Calorimetry experiment ('XQC', excluded region) and halo heating and stability arguments ('Halo', excluded above) provide model-independent limits. Under appropriate assumptions about the behavior of dark matter in the early universe, the Cosmic Microwave Background ('CMB', excluded above) provides a tighter bound for $m_{\text{DM}} < 1$ MeV. This can be slightly improved ('CMB+') with arguments from Large Scale Structure data, elemental abundances from big bang nucleosynthesis, and the Hubble constant H_0. On top of these existing constraints, the possible sensitivities of four proposed experiments are shown: magnetically trapped superconducting microspheres [47] ('Pb sphere'), optically-trapped silican nanospheres [48] ('Si sphere'), a laser-driven silican mirror [49] ('Si mirror'), and the optical time-domain ionizing matter-wave interferometer [50] ('OTIMA') for several choices of cluster mass. Significant visibility boosts for all experiments exist below roughly 1 MeV, where the typical de Broglie wavelength of the dark matter is long enough to scatter coherently off multiple nuclei. (Coherent scattering off multiple nucleons in the same nucleus occurs for the entire plotted range.) When the de Broglie wavelength is larger than the size of atoms but smaller than the size of the interfered object, complicated scattering effects exist. The dashed lines on the experiments are regions where a very crude fractional coherence boosts approximation has been made. Below roughly 10 eV, the number density of cold dark matter becomes so large that Fermi degeneracy prevents fermionic dark matter from reaching the density necessary to explain galactic rotation curves. Strongly interacting massive particle ('SIMP') models probably cannot be accessed by decoherence-based approaches in the near future, but it provides a useful reference.

scattering $|f(q\hat{q}, q\hat{q}')|^2 \propto 1$, $\sigma^{\rm tot}$ and $\sigma^{\rm tr}$ coincide. If we assume $\sigma \equiv \sigma^{\rm tot} = \sigma^{\rm tr}$ varies slowly with q, we can then numerically compute the integrals in Eq. 17.

Note that the decoherence factor γ is multiplicative for independent scattering events, so for a time-varying separation $\Delta x(t)$ in the interferometer, one can simply use the time average $\langle \Delta x^2 \rangle = T^{-1} \int_0^T \Delta x^2(t)\,\mathrm{d}t$ when computing γ. The condition for effective decoherence is that $\gamma \ll 1$, that is $T \gtrsim 1/F(\Delta x)$.

Interferometry with single atoms has little hope of being sensitive to dark matter because of the rarity of interactions. Fortunately, recent progress in the size of superposed objects in matter-wave interferometry has been stunning, with clear fringe patterns produced when interfering molecules composed of up to 430 atoms and in excess of 6,000 amu [56]. Optical time-domain ionizing matter-wave (OTIMA) interferometers proposed by Nimmrichter, Haslinger, Hornberger, and Arndt (NHHA) will interfere clusters of matter larger than 10^6 amu [50]. For the purpose of demonstrating the potential of dark matter detection through decoherence-based techniques, I will use the NHHA proposal as a benchmark and roughly estimate its sensitivity to dark matter for different cluster sizes. See Fig. 2.

Modern matter-wave interferometry experiments like the NHHA proposal use multiple gratings with many slits to overcome difficulties with beam coherence and tiny de Broglie wavelengths [29, 30], so the matter clusters are not described by a simple superposition of two spatially separated wavepackets. Nonetheless, treating the experiment this way should gives us a good estimate of the potential visibility of dark matter. For this purpose, we choose $\Delta x = 157$ nm to be equal to the wavelength of the laser grating used to produce the "slits" in the NHHA proposal. More detailed calculations are deferred to future studies. Note that only the results for $m_{\rm DM} \lesssim 20$ keV will depend on this choice. For larger masses, which are in the short wavelength limit, any scattering event results in complete decoherence independent of Δx.

Importantly, sub-GeV dark matter will scatter coherently off multiple nucleons so long as its coupling to nucleons is independent of spin, as is commonly assumed. This means that the total scattering rate is proportional to the *square* of the number of nucleons in the target so long as the target is smaller than the de Broglie wavelength of the dark matter, i.e. the target fits inside the coherence volume λ^3. When the wavelength is large enough to span multiple nuclei, but not large enough to span the whole target, complicated interference effects (both constructive and destructive) arise which depend sensitively on the angular cross section and the distribution of nuclei. For the estimates in this work, I make the crude assumption that the boost is linear in the number of nuclei which fit in the coherence volume, deferring more accurate calculations.

In the same way, I also estimate the visibility of three other proposed experiments: magnetically trapped superconducting microspheres [47], optically-trapped silican nanospheres [48], and a photon-driven oscillating mirror [49].

Direct-detection experiments on the Earth's surface will only be sensitive to dark matter if its cross-section with nucleons is sufficiently low for it to pass through the atmosphere and reach the experiment. This upper bounds the spin-independent cross-section visible to Earth-bound experiments at about $10^{-27}\mathrm{cm}^2$. This can be circumvented by placing the experiment on a sounding balloon (~30 km altitude; $10^{-25}\mathrm{cm}^2$), a sounding rocket (~200 km altitude; $10^{-18}\mathrm{cm}^2$), or on a satellite in low-Earth orbit. I do not know if anyone has studied the possibility of observing non-classical states on a balloon or rocket, but spaceborne experiments are certainly feasible [50, 57, 58]. As shown in Fig. 2, the next generation of quantum superposition experiments would probably need to be raised at least partially out of the Earth's atmosphere to see dark matter.

The only existing direct-detection constraints on the spin-independent dark-matter-nucleon scattering cross-section σ for masses $m_{\rm DM}$ below a GeV come from the X-ray Quantum Calorimetry (XQC) experiment [59], as analyzed by Erickcek et al. [60]. (See, however, [61].) There is additionally a pair of related model-independent bounds arising from the stability of dark matter halo and the rate at which it heats interstellar gas through collisions [62].

Although it's tempting to remain agnostic about the behavior of the dark sector in the early universe, the success of cosmological models in explaining observations allow additional constraints with some confidence. Assuming the standard thermal freeze-out scenario, the Cosmic Microwave Background (CMB) puts tighter upper bounds on σ than those from halo heating and stability for $m_{\rm DM} > 1$ MeV [63]. They can slightly improved by the addition of constraints arising from Large Scale Structure data (LSS), elemental abundances from big bang nucleosynthesis (BBN), and the Hubble constant H_0.

These contraints are all plotted in Fig. 2. Without making significant new assumptions, there does not appear to be any other restrictions on cold dark matter for low masses.

As stated in the previous section, the many possible sources of decoherence mean that its mere detection has ambiguous implication. The observations of interference (i.e. the elimination of all decoherence) can reliably exclude dark matter parameter space, but the mere detection of decoherence is not enough to conclusively demonstrate the existence of dark matter particles. To do that, experimenters must have some handle on the degree of dark-matter originating decoherence. By manipulating the expected amount of decoherence, and confirming that it matches what is observed in the experiment, one can establish enough properties of the source to be confident the source is dark matter.

If the dark matter interacts strongly enough to be blocked by normal matter, then moving the experiment underground would reduce decoherence and increase fringe visibility, while placing the experiment above the atmosphere would increase decoherence and reduce fringe visibility. On the other hand, if normal matter is ineffective at shielding the dark matter, the orbit of the earth around the sun could be used to see an annual modulation. (See, for example, the Michelson-Morely experiment [64, 65], and the CoGeNT [1] and DAMA family [3, 66, 67] of dark matter searches.) When the earth is moving with the stream of new particles, flux and decoherence would be less; when the Earth is moving against the stream, flux and decoherence would be greater.

Likewise, if interferometers were used to detect hypothetical particles produced artificially on earth—dark matter or otherwise—then the velocity of the hypothetical particles could be slow enough that physically moving the interferometer in the laboratory modulates the signal.

Gravitons, fifth forces, and relic neutrinos

Gravitons are generally thought to be undetectable by any feasible measurement [68–70]. In particular, a Jupiter-sized detector in close orbit around a neutron star is expected to absorb no more than one graviton every decade, and even then is unlikely to be able to distinguish it from background events in any imaginable manner [68]. However, a non-classical detection scheme may be able to reveal gravitons—although the technology for such an experiment still lies far in the future.

Consider an experiment where a clump of matter of mass m is brought into a superposition of two spatially localized center-of-mass wavepackets separated by a distance L and then recombined after a time τ, so that the interference pattern can be observed. The object accelerates to a speed $v = \beta c \sim L/\tau$. See Fig. 3. Even if there is no decoherence of the two paths from external environments, the possibility of *intrinsic* decoherence due to emitted electromagnetic radiation is well known [71, 72]. Assuming the clump is thermal, there are two types of possible decohering radiation: blackbody and bremsstrahlung. Decoherence from blackbody radiation can be avoided by cooling the body to a temperature with characteristic wavelength $\lambda_T \gg L$.

Breuer and Pettruccione [71, 73] calculate the decoherence due to electromagnetic bremsstrahlung for this experiment and found that interference between the two paths is suppressed by a decoherence factor γ where

$$\ln\gamma \approx -\frac{2\alpha_e}{\pi}C_0\beta^2 \tag{18}$$

for negligible temperature (initial vacuum state) of the electromagnetic field in the non-relativistic ($\beta \ll 1$) limit. Above, $\alpha_e = q^2/4\pi\hbar c$ is the effective electromagnetic coupling constant for a clump of charge q and C_0 is a constant of order unity which depends only logarithmically on an infrared cutoff set by the finite duration τ of the superposition. In this non-relativistic limit, the photons have wavelength of order $\lambda = \beta^{-1}L \gg L$ and it takes many photons to decohere. For relativistic β, the wavelength of the radiation is comparable to L. In this case, it turns out that the decoherence factor γ is essentially the probability that the clump does not emit any photons, which still goes like $\ln\gamma \sim -\alpha_e\beta^2$. Notice that this does not strongly depend on L so long as L is larger than the width of the wavepackets.



FIG. 3. Decoherence by electromagnetic or gravitational bremsstrahlung. A charged object traveling at a speed is brought into a coherent superposition of spatial extent L then recombined after a time τ. If the two paths include sufficient relative acceleration for the given charge, they decohere through the emission of bremsstrahlung, which records with-path information. The radiation need not be absorbed for its existence to be detected.

Electromagnetic bremsstrahlung can be avoided by ensuring that there is no net electromagnetic charge. (Microscopic charge inhomogeneities are smoothed out on scales smaller than λ.) The same is not true for *gravitational* bremsstrahlung because the mass of the object is always non-zero. We cannot directly use the results of Breuer and Pettruccione because gravitons are spin-2 and gravitational radiation does not have a dipole component. However, from the calculation of electromagnetic and gravitational quadrupole radiation [74, 75], which picks up an extra factor of β^2, it's clear that the decoherence factor behaves like

$$\ln\gamma \sim -\alpha_g\beta^4 = -\frac{Gm^2}{\hbar c}\beta^4. \tag{19}$$

The planck mass

$$m_P = \sqrt{\frac{\hbar c}{G}} \approx 1.2\times 10^{19}\mathrm{GeV}/c^2 \approx 21\mu\mathrm{g} \tag{20}$$

is precisely the mass scale at which the gravitational coupling constant α_g reaches unity. Thus, the coherent manipulation of Planck-mass superpositions at relativistic speeds will enable the detection of gravitons. Non-relativistic speeds necessitate an increase in mass by a factor β^{-2}.

As mentioned, matter-wave interferometry has been demonstrated with $m \sim 10^4$ amu, with no fundamental obstacles to surpassing $m \sim 10^6$ amu [50]. The primary barrier to experiments interfering masses larger than $m \sim 10^7$ amu is the Earth's gravity, which may be eliminated in spaceborne experiments [50, 76]. Superpositions of lead spheres with $m \sim 10^{14}$ amu [47, 48] and of oscillating mirrors with $m \sim 10^{16}$ amu [49, 77] are being pursued, although the spatial extents of such superpositions are too small to decohere through bremsstrahlung. Given this, the coherent manipulation of Planck-mass (10^{19} amu) objects—though massively difficult—is not inconceivable. It is arguably more feasible than constructing a detector of Jovian proportions.

The key aspect of this experiment is that the elementary masses making up the lump of matter act together coherently to generate the graviton. Note that the emitted graviton *cannot* be observed by any feasible classical detection scheme. Nonetheless, its presence can be inferred by the decoherence it causes, just like the presence of a decohering gas in existing interferometers.

Of course, such a scheme for detecting the presence of gravitons assumes that all other sources of decoherence can be suppressed. There are at least two such sources which, if they hinder the observation of gravitons, would themselves be exciting new physics: collisional decoherence from relic neutrinos, and bremsstrahlung from fifth forces stronger than gravity. The investigation of these possibilities is deferred to future work.

Relation to standard quantum limits

A *standard quantum limit* (SQL) refers most generally to a limit on experimental accuracy due to quantum effects. It appears that all SQLs existing in the literature can be formulated in terms of a classical detection scheme, that is, the estimation of a parameter through the preparation and measurement of a quantum system in a single given basis. It is satisfying to explicitly reproduce two of the most well known SQLs: the accuracy with which the displacement of test masses can be measured (often known as *the* standard quantum limit) and the shot noise limit for probing local unitaries with unentangled photons. When placed in this framework, its clear that existing SQLs are distinct from the decoherence-based technique introduced in this article.

For concreteness I will use the classical fidelity (Bhattacharyya distance)

$$F(p,q) = \sum_i \sqrt{p_i q_i} \tag{21}$$

as a measure of the distinguishability of two probability distributions p_i and q_i. Note that $F = 1$ if $p_i = q_i$ and $F = 0$ if p_i and q_i are perfectly distinguishable, i.e. if their supports are disjoint.

First, consider a weak force which operates on an otherwise free particle for a time T between an initial preparation and a (POVM) measurement. Suppose we are restricted to a classical detection scheme in the overcomplete basis of Gaussian wavepackets of spatial width σ_0 and zero expected momentum. A wavepacket under the influence of a classical force has a center in phase space which follows the classical equations of motion and (regardless of whether the classical force is applied) has a spatial width which grows like [78]

$$\sigma(T)^2 = \sigma_0^2 + (\hbar T/2m\sigma_0)^2. \tag{22}$$

The fidelity between the probability distributions for the result of the POVM measurement, with and without the force, is $F = \exp[-\Delta x_F^2/8(\sigma(T)^2+\sigma_0^2)]$, where Δx_F is the classical displacement caused by the force. The fidelity is minimized (distinguishability maximized) when $\sigma_0^2 = \hbar T/2\sqrt{2}m$. (In some cases, the effective value of σ_0 may be controllable by adjusting macroscopic parameters of the environment [79].) This motivates defining the standard quantum limit

$$\Delta x_{\mathrm{SQL}}^2 = 2\sqrt{2}\hbar T/m. \tag{23}$$

Since

$$F = e^{-\Delta x_F^2/4\Delta x_{\mathrm{SQL}}^2} \tag{24}$$

the force is only detectable when $\Delta x_F \gtrsim \Delta x_{\mathrm{SQL}}$.

Naturally, this SQL for test masses can be beaten by preparing non-classical superpositions. For classical displacements much smaller than the SQL, $\Delta x_F \ll \Delta x_{\mathrm{SQL}}$, the effect of the force is to add a phase $\chi = \Delta x p_0/\hbar$ to the wavepacket (with negligible effect on its spatial envelope), where p_0 is the momentum of the test mass. This phase can be observed by preparing a coherent spatial superposition of wavepackets, such as using atom interferometry to detect gravitational waves [80].

Second, consider N qubits with a Hilbert space of dimension 2^N. The qubits are acted upon by local unitaries which are all identically equal to V_a or V_b, which are nearby: $V_b = UV_a$ for U near the identity. Since an overall phase difference between V_a and V_b cannot be observed, we can set one of the eigenvalues of U to unity and let the computation basis diagonalize it:

$$U = |0\rangle\langle 0| + e^{i\phi}|1\rangle\langle 1|. \tag{25}$$

The goal is to optimally distinguish between V_a and V_b for small ϕ. For a single qubit ($N = 1$), an arbitrary initial state and measurement basis leads to two probability distributions for the outcomes. The fidelity between them is

$$F = 1 - \chi\phi^2 + O(\phi^3), \tag{26}$$

where $\chi \leq 1/4$ is a non-negative constant depending on the initial state and the measurement basis. For N *unentangled* qubits measured in an arbitrary product basis, the fidelity is

$$F = \prod_{i=1}^{N} F_i = 1 - \langle\chi\rangle N\phi^2 + O(\phi^3). \tag{27}$$

This is the shot noise limit for detecting a small phase in a classical detection scheme utilizing an unentangled basis for preparation and measurement: for the unitaries to be distinguishable for large N, ϕ must decrease no faster than $1/\sqrt{N}$.

On the other hand, if we allow the initial entangled state

$$|\psi_{\mathrm{in}}\rangle = (V_a^{\otimes N})^{-1}\frac{|0\rangle^{\otimes N} + |1\rangle^{\otimes N}}{\sqrt{2}}, \tag{28}$$

we can achieve an accuracy in ϕ which scales like $1/N$. This is the so-called Heisenberg limit, and it is the best one can do [81–83].

These examples, and all SQL's of which I am aware, assume *unitary* dynamics of the measured system. In other words, they treat the environment completely classically. (The "decoherence-enhanced measurement" of Braun and Martin [84]—though a promising idea for evading the shot-noise limit in a decoherence-resistant manner—is misleadingly named. Under that protocol, the environment responsible for the decoherence actually serves as an auxiliary system and must be measured.) Furthermore, if we move to the exact "classical limit", such as $\Delta x_F = 0$ or $\phi = 0$, the phenomena become completely undetectable, whether or not quantum methods are employed.

In contrast, the decoherence-based schemes proposed in this article are crucially non-unitary; the dynamics increase the mixedness of the measured system. Most strikingly, they can enable detection of phenomena which, classically, cannot be detected by *any* means. Hypothetical particles transferring exactly zero momentum can still easily cause decoherence in a matter-wave interferometer.

DISCUSSION

One can reinterpret many experiments which establish non-classical states as direct evidence against hypothetical phenomena which, if they existed, would decohere those states. Insofar as stability in the presense of decoherence defines the degree of classicality of a state (an extension of the fact that *the* classical pointer states are the *most* robust to decoherence [10, 11, 15]), the best constraints on hypothetical weak phenomena will come from the most non-classical states.

It's not clear whether any of the particular proposals I have discussed will turn out to be used for detecting new phenomena. Evolution is usually not exactly diagonal in the pointer basis, so any given proposal might be surpassed by a sufficiently sensitive classical detection scheme. Nonetheless, the amazing properties of non-classical schemes gives good reasons to investigate them. In addition, many experiments investigating non-classical states are already being pursued with completely independent motivations; the ability of these experiments to rule out weak phenomena (or discover them!) is an added bonus.

Acknowledgements

I thank Asimina Arvanitaki, Dirk Bouwmeester, Savas Dimopoulos, Rouven Essig, Alexander Friedland, Christian Hagmann, Andrew Hime, Lorenzo Maccone, Jeremy Mardon, Harry Nelson, Bryon Neufeld, Shmuel Nussinov, Keith Rielage, Daniel Sank, Alexia Schulz, Mark Srednicki, Paul Steinhardt, and Haibo Yu for discussion. I especially thank Michael Graesser for patient teaching, Godfrey Miller for thoughtful criticism, and Wojciech Zurek and Jim Hartle for making this work possible. This research was partially supported by the U.S. Department of Energy through the LANL/LDRD program.

---

[1] C. E. Aalseth, P. S. Barbeau, J. Colaresi, J. I. Collar, J. Diaz Leon, J. E. Fast, N. Fields, T. W. Hossbach, M. E. Keillor, J. D. Kephart, A. Knecht, M. G. Marino, H. S. Miley, M. L. Miller, J. L. Orrell, D. C. Radford, J. F. Wilkerson, and K. M. Yocum (The CoGeNT Collaboration), Phys. Rev. Lett., **107**, 141301 (2011).

[2] M. Bravin, M. Bruckmayer, C. Bucci, S. Cooper, S. Giordano, F. von Feilitzsch, J. Hhne, J. Jochum, V. Jrgens, R. Keeling, H. Kraus, M. Loidl, J. Lush, J. Macallister, J. Marchese, O. Meier, P. Meunier, U. Nagel, T. Nssle, F. Prbst, Y. Ramachers, M. Sarsa, J. Schnagl, W. Seidel, I. Sergeyev, M. Sisti, L. Stodolsky, S. Uchaikin, and L. Zerle (The CRESST Collaboration), Astroparticle Physics, **12**, 107 (1999), ISSN 0927-6505.

[3] R. Bernabei, P. Belli, A. Bussolotti, F. Cappella, R. Cerulli, C. J. Dai, A. D'Angelo, H. L. He, A. Incicchitti, H. H. Kuang, J. M. Ma, A. Mattei, F. Montecchia, F. Nozzoli, D. Prosperi, X. D. Sheng, and Z. P. Ye, Nuclear Instruments and Methods in Physics Research A, **592**, 297 (2008), arXiv:0804.2738.

[4] Z. Ahmed, D. S. Akerib, S. Arrenberg, M. J. Attisha, C. N. Bailey, L. Baudis, D. A. Bauer, J. Beaty, P. L. Brink, T. Bruch, R. Bunker, S. Burke, B. Cabrera, D. O. Caldwell, J. Cooley, P. Cushman, F. DeJongh, M. R. Dragowsky, L. Duong, J. Emes, E. Figueroa-Feliciano, J. Filippini, M. Fritts, R. J. Gaitskell, S. R. Golwala, D. R. Grant, J. Hall, R. Hennings-Yeomans, S. Hertel, D. Holmgren, M. E. Huber, R. Mahapatra, V. Mandic, K. A. McCarthy, N. Mirabolfathi, H. Nelson, L. Novak, R. W. Ogburn, M. Pyle, X. Qiu, E. Ramberg, W. Rau, A. Reisetter, T. Saab, B. Sadoulet, J. Sander,

R. Schmitt, R. W. Schnee, D. N. Seitz, B. Serfass, A. Sirois, K. M. Sundqvist, M. Tarka, A. Tomada, G. Wang, S. Yellin, J. Yoo, and B. A. Young (The CDMS Collaboration), Science, **327**, 1619 (2010), http://www.sciencemag.org/content/327/5973/1619.full.pdf.

[5] E. Aprile, K. Arisaka, F. Arneodo, A. Askin, L. Baudis, A. Behrens, K. Bokeloh, E. Brown, J. M. R. Cardoso, B. Choi, D. B. Cline, S. Fattori, A. D. Ferella, K.-L. Giboni, A. Kish, C. W. Lam, J. Lamblin, R. F. Lang, K. E. Lim, J. A. M. Lopes, T. Marrodán Undagoitia, Y. Mei, A. J. Melgarejo Fernandez, K. Ni, U. Oberlack, S. E. A. Orrigo, E. Pantic, G. Plante, A. C. C. Ribeiro, R. Santorelli, J. M. F. dos Santos, M. Schumann, P. Shagin, A. Teymourian, D. Thers, E. Tziaferi, H. Wang, and C. Weinheimer (The XENON100 Collaboration), Phys. Rev. Lett., **105**, 131302 (2010).

[6] J. L. Feng, J.-F. m. c. Grivaz, and J. Nachtman, Rev. Mod. Phys., **82**, 699 (2010).

[7] Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, **599**, 28 (2009), ISSN 0168-9002.

[8] S. N. Gninenko, International Journal of Modern Physics A, **19**, 3833 (2004).

[9] E. Fischbach and C. Talmadge, Nature, **356**, 207 (1992).

[10] W. H. Zurek, Phys. Rev. D, **24**, 1516 (1981).

[11] W. H. Zurek, Phys. Rev. D, **26**, 1862 (1982).

[12] W. H. Zurek, S. Habib, and J. P. Paz, Phys. Rev. Lett., **70**, 1187 (1993).

[13] M. Schlosshauer, *Decoherence and the Quantum-to-Classical Transition* (Springer-Verlag, Berlin, 2008).

[14] E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu, *Decoherence and the Appearance of the Classical World in Quantum Theory*, 2nd ed. (Springer-Verlag, Berlin, 2003).

[15] W. H. Zurek, Rev. Mod. Phys., **75**, 715 (2003).

[16] H. Ollivier, D. Poulin, and W. H. Zurek, Phys. Rev. A, **72**, 042113 (2005).

[17] W. H. Zurek, Nature Physics, **5**, 181 (2009).

[18] H. Ollivier, D. Poulin, and W. H. Zurek, Phys. Rev. Lett., **93**, 220401 (2004).

[19] H. O'Keeffe, E. OSullivan, and M. Chen, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, **640**, 119 (2011), ISSN 0168-9002.

[20] Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, **449**, 172 (2000), ISSN 0168-9002.

[21] E. Aprile and T. Doke, Rev. Mod. Phys., **82**, 2053 (2010).

[22] W. Ehrenberg and R. E. Siday, Proceedings of the Physical Society. Section B, **62**, 8 (1949).

[23] Y. Aharonov and D. Bohm, Phys. Rev., **115**, 485 (1959).

[24] R. G. Chambers, Phys. Rev. Lett., **5**, 3 (1960).

[25] G. Möllenstedt and W. Bayh, Naturwissenschaften, **49**, 81 (1962), ISSN 0028-1042, 10.1007/BF00622023.

[26] A. Tonomura, N. Osakabe, T. Matsuda, T. Kawasaki, J. Endo, S. Yano, and H. Yamada, Phys. Rev. Lett., **56**, 792 (1986).

[27] G. d. f. Matteucci and G. Pozzi, Phys. Rev. Lett., **54**, 2469 (1985).

[28] L. Vaidman, pre-print (2011), arXiv:1110.6169.

[29] A. D. Cronin, J. Schmiedmayer, and D. E. Pritchard, Rev. Mod. Phys., **81**, 1051 (2009).

[30] K. Hornberger, S. Gerlich, P. Haslinger, S. Nimmrichter, and M. Arndt, Review of Modern Physics, *in press* (2011), arXiv:1109.5937.

[31] K. Hornberger, S. Uttenthaler, B. Brezger, L. Hackermüller, M. Arndt, and A. Zeilinger, Phys. Rev. Lett., **90**, 160401 (2003).

[32] H. Uys, J. D. Perreault, and A. D. Cronin, Phys. Rev. Lett., **95**, 150403 (2005).

[33] S. Nimmrichter, K. Hornberger, P. Haslinger, and M. Arndt, Phys. Rev. A, **83**, 043621 (2011).

[34] G. C. Ghirardi, A. Rimini, and T. Weber, Phys. Rev. D, **34**, 470 (1986).

[35] B. Vacchini, Journal of Physics A: Mathematical and Theoretical, **40**, 2463 (2007).

[36] R. Catena and P. Ullio, Journal of Cosmology and Astroparticle Physics, **2010**, 004 (2010).

[37] J. Lewin and P. Smith, Astroparticle Physics, **6**, 87 (1996), ISSN 0927-6505.

[38] S. L. Adler, Phys. Rev. D, **79**, 023505 (2009).

[39] B. W. Lee and S. Weinberg, Phys. Rev. Lett., **39**, 165 (1977).

[40] S. Dodelson and L. M. Widrow, Phys. Rev. Lett., **72**, 17 (1994).

[41] K. Abazajian, G. M. Fuller, and M. Patel, Phys. Rev. D, **64**, 023501 (2001).

[42] K. Abazajian, Phys. Rev. D, **73**, 063506 (2006).

[43] C. T. Kishimoto and G. M. Fuller, Phys. Rev. D, **78**, 023524 (2008).

[44] M. Pospelov, A. Ritz, and M. Voloshin, Phys. Rev. D, **78**, 115012 (2008).

[45] K. M. Z. Tongyan Lin, Hai-Bo Yu, (2011), arXiv:1111.0293.

[46] R. Essig, J. Mardon, and T. Volansky, (2012), arXiv:1108.5383.

[47] O. Romero-Isart, L. Clemente, C. Navau, A. Sanchez, and J. I. Cirac, (2012), arXiv:1112.5609.

[48] O. Romero-Isart, A. C. Pflanzer, F. Blaser, R. Kaltenbaek, N. Kiesel, M. Aspelmeyer, and J. I. Cirac, Phys. Rev. Lett., **107**, 020405 (2011).

[49] W. Marshall, C. Simon, R. Penrose, and D. Bouwmeester, Phys. Rev. Lett., **91**, 130401 (2003).

[50] S. Nimmrichter, P. Haslinger, K. Hornberger, and M. Arndt, New Journal of Physics, **13**, 075002 (2011).

[51] E. Joos and H. D. Zeh, Zeitschrift für Physik B Condensed Matter, **59**, 223 (1985).

[52] M. R. Gallis and G. N. Fleming, Phys. Rev. A, **42**, 38 (1990).

[53] L. Diosi, Europhys. Lett., **30**, 63 (1995).

[54] K. Hornberger and J. E. Sipe, Phys. Rev. A, **68**, 012105 (2003).

[55] K. Hornberger, Phys. Rev. Lett., **97**, 060601 (2006).

[56] S. Gerlich, S. Eibenberger, M. Tomandl, S. Nimmrichter, K. Hornberger, P. J. Fagan, J. Tüxen, M. Mayor, and M. Arndt, Nat Commun, **2** (2011).

[57] R. Kaltenbaek, G. Hechenblaikner, N. Kiesel, U. Johann, and M. Aspelmeyer, in *Lasers and Electro-Optics Europe (CLEO EUROPE/EQEC), 2011 Conference on and 12th European Quantum Electronics Conference* (2011) p. 1.

[58] R. Kaltenbaek, G. Hechenblaikner, N. Kiesel, O. Romero-Isart, U. Johann, and M. Aspelmeyer, (2012), arXiv:1201.4756.

[59] D. McCammon, R. Almy, E. Apodaca, W. B. Tiest, W. Cui, S. Deiker, M. Galeazzi, M. Juda, A. Lesser, T. Mihara, J. P. Morgenthaler, W. T. Sanders, J. Zhang,

E. Figueroa-Feliciano, R. L. Kelley, S. H. Moseley, R. F. Mushotzky, F. S. Porter, C. K. Stahle, and A. E. Szymkowiak, The Astrophysical Journal, **576**, 188 (2002).

[60] A. L. Erickcek, P. J. Steinhardt, D. McCammon, and P. C. McGuire, Phys. Rev. D, **76**, 042007 (2007).

[61] J. P. S. . T. V. Rouven Essig, Aaron Manalaysay, forthcoming.

[62] R. S. Chivukula, A. G. Cohen, S. Dimopoulos, and T. P. Walker, Phys. Rev. Lett., **65**, 957 (1990).

[63] X. Chen, S. Hannestad, and R. J. Scherrer, Phys. Rev. D, **65**, 123515 (2002).

[64] A. A. Michelson, American Journal of Science, **22**, 120 (1881).

[65] A. A. Michelson and E. W. Morley, American Journal of Science, **34**, 333 (1887).

[66] R. Bernabei, P. Belli, F. Cappella, R. Cerulli, F. Montechia, F. Nozzoli, A. Incicchitti, D. Prosperi, C. J. Dai, H. H. Kuang, J. M. Ma, and Z. P. Ye, Nuovo Cimento Rivista Serie, **26**, 010000 (2003), arXiv:astro-ph/0307403.

[67] J. Cherwinka, R. Co, D. F. Cowen, D. Grant, F. Halzen, K. M. Heeger, L. Hsu, A. Karle, V. A. Kudryavtsev, R. Maruyama, W. Pettus, M. Robinson, and N. J. C. Spooner, (2011), arXiv:1106.1156.

[68] T. Rothman and S. Boughn, Foundations of Physics, **36**, 1801 (2006), ISSN 0015-9018, 10.1007/s10701-006-9081-9.

[69] S. Boughn and T. Rothman, Classical and Quantum Gravity, **23**, 5839 (2006).

[70] F. J. Dyson, in *NY Rev. Books 51 (8)*, edited by M. Kafatos (2004).

[71] H.-P. Breuer and F. Petruccione, Phys. Rev. A, **63**, 032102 (2001).

[72] D. Drr and H. Spohn, in *Decoherence: Theoretical, Experimental, and Conceptual Problems*, Lecture Notes in Physics, Vol. 538, edited by P. Blanchard, E. Joos, D. Giulini, C. Kiefer, and I.-O. Stamatescu (Springer Berlin / Heidelberg, 2000) pp. 77–86, ISBN 978-3-540-66899-2.

[73] H.-P. Breuer and F. Petruccione, "The theory of open quantum systems," (Oxford University Press, USA, 2002) pp. 577–605.

[74] R. J. Gould, The Astrophysical Journal, **288**, 789 (1985).

[75] S. Weinberg, Phys. Rev., **140**, B516 (1965).

[76] M. Arndt, (2012), private communication.

[77] D. Vitali, S. Gigan, A. Ferreira, H. R. Böhm, P. Tombesi, A. Guerreiro, V. Vedral, A. Zeilinger, and M. Aspelmeyer, Phys. Rev. Lett., **98**, 030405 (2007).

[78] J. R. Choi, M. Maamache, Y. Saadi, and K. H. Yeon, Journal of the Korean Physical Society, **56**, 1063 (2010).

[79] L. Diósi and C. Kiefer, Phys. Rev. Lett., **85**, 3552 (2000).

[80] S. Dimopoulos, P. W. Graham, J. M. Hogan, M. A. Kasevich, and S. Rajendran, Phys. Rev. D, **78**, 122002 (2008).

[81] Z. Y. Ou, Phys. Rev. A, **55**, 2598 (1997).

[82] V. Giovannetti, S. Lloyd, and L. Maccone, Phys. Rev. Lett., **96**, 010401 (2006).

[83] V. Giovannetti, S. Lloyd, and L. Maccone, Nat Photon, **5**, 222 (2011).

[84] D. Braun and J. Martin, Nature Communications, **2**, 223 (2011).